White Shark Captures by Season in Southern California

Spring




Figure 14.3 Spatial distribution of reported White Shark captures occurring in Southern California during spring (April to June, a), summer (July to September, b), fall (October to December, c), and winter (January to March, d), 1935-2009. The numbers represent sample sizes for fishing blocks with greater than 10 sharks.

Copyright 2012. From *Global Perspectives on the Biology and Life History of the White Shark*, Chapter 14 "Historic Fishery Interactions with White Sharks in the Southern California Bight" by C. Lowe, M. Blasius, E. Jarvis, T. Mason, G. Goodman-Lowe and J. O'Sullivan. Reproduced by permission of Taylor and Francis Group, LLC, a division of Informa plc.

A project of CA NGSS K-8 Early Implementation Initiative.



Toolbox 8.9

Group Handout

8.9.G2



White Shark Captures by Season in Southern California (continued)

Summer




Figure 14.3 Spatial distribution of reported White Shark captures occurring in Southern California during spring (April to June, a), summer (July to September, b), fall (October to December, c), and winter (January to March, d), 1935-2009. The numbers represent sample sizes for fishing blocks with greater than 10 sharks.

A project of CA NGSS K-8 Early Implementation Initiative.





White Shark Captures by Season in Southern California (continued)

Fall




Figure 14.3 Spatial distribution of reported White Shark captures occurring in Southern California during spring (April to June, a), summer (July to September, b), fall (October to December, c), and winter (January to March, d), 1935-2009. The numbers represent sample sizes for fishing blocks with greater than 10 sharks.

A project of CA NGSS K–8 Early Implementation Initiative.




Figure 14.3 Spatial distribution of reported White Shark captures occurring in Southern California during spring (April to June, a), summer (July to September, b), fall (October to December, c), and winter (January to March, d), 1935-2009. The numbers represent sample sizes for fishing blocks with greater than 10 sharks.

A project of CA NGSS K–8 Early Implementation Initiative.